



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

to 14<sup>h</sup> 55<sup>m</sup> fifty-one meteors were counted, of which all but four were *Perseids*. This frequency was estimated to be about an average for the latter portion of the night. C. D. PERRINE.

LICK OBSERVATORY,  
UNIVERSITY OF CALIFORNIA, August 16, 1898.

ELEMENTS OF COMET *e* 1898 (PERRINE).

The following system of parabolic elements of this comet has been derived from normal places for the dates June 16.0, July 12.0, and August 7.0. The observations used in forming the normal places were: Mount Hamilton, June 14, 15, 16, 17; Paris, June 16; Strassburg, June 17; Mount Hamilton, July 9, 11, 12, 13, 14; Mount Hamilton, August 2, 4, 5, 6, 7, 8.

ELEMENTS.

$T = 1898$ , August 16.19978 G. M. T.

$$\left. \begin{array}{lll} \Omega = 259^\circ & 6' & 12''.2 \\ \omega = 205 & 36 & 24.0 \\ i = 70 & 1 & 36.7 \end{array} \right\} \begin{array}{l} \text{Ecliptic and} \\ \text{Mean Equinox 1898.0} \end{array}$$

$$\log q = 9.796950$$

The residuals for the middle place are:—

Observed — Computed,

$$\begin{array}{ll} \Delta \lambda' \cos \beta' & + 0''.1 \\ \Delta \beta' & - 0.9 \end{array}$$

The comet was last observed at Mount Hamilton on the morning of August 11th, when it was well into the dawn. It would not have been visible except for its increased brightness and sharp nucleus. On August 7th, the nucleus of the comet was estimated to be nearly as bright as the 9.1-magnitude comparison star. The light of the entire comet probably equaled a seventh-magnitude star. The comet has now passed out of range of northern observatories, but should be visible in the Southern Hemisphere for two months yet. The orbit of this comet bears a resemblance to that of the PONS-BROOKS comet of 1812–1884. There is also considerable resemblance to the orbit of the comet 1785 I, especially in  $\omega$  and  $i$ . Comet *e* is so plainly parabolic, that the resemblance must be considered as merely placing them in a group, probably with no physical connection. C. D. PERRINE.

LICK OBSERVATORY,  
UNIVERSITY OF CALIFORNIA, August 25, 1898.